

Macrophage Functions after Exposure to Mineral Fibers

by F. Tilkes* and E. G. Beck*

UICC, other well-defined asbestos samples and different man-made mineral fibers (MMM) such as glass fiber and synthetic amphibole asbestos were studied *in vitro* by using rat and guinea pig lung macrophages. These samples had relatively narrow length and diameter spectra.

Most of the fiber samples were added to the cultures on a gravimetric basis, although some were added on a numerical basis. Electrocorundum and DQ₁₂ (Dorentruper Quartz) were used as controls at comparable gravimetric concentrations.

The assays used were the release of lactate dehydrogenase (to demonstrate plasma membrane permeability) and the release of β -glucuronidase (to indicate lysosomal permeability). Carbohydrate metabolism was monitored by the measurement of lactic acid production and, as one of the tests for macrophage function, the production of lysozyme was determined.

The phagocytic ability of the cells was measured, after the addition of opsonized zymosan, by bioluminescence following luminol enhancement. Only some results could be evaluated, however, due to technical difficulties.

A length- and dose-dependent cytotoxicity of the fibers was found in this system which was similar to that previously described with permanent cell lines. No great differences were found between fibers having different physicochemical compositions if their geometric dimensions were similar. Long, very thin fibers of glass, chrysotile, crocidolite and synthetic fluoroamphiboles were all toxic in the test system.

Introduction

The macrophage plays a central role in the manifestation of inflammation (1). Furthermore, macrophages are the first target cells with which environmental and occupational dusts come into contact after they have been inhaled into the lung (2).

Silica, and asbestos fibers, have been recognized as being toxic in *in vitro* macrophage cell systems and this cytotoxicity is characterized by the release of cytoplasmic and lysosomal enzymes (3, 4). However, Davies et al. (5) have demonstrated that peritoneal macrophages can also release acid lysosomal hydrolases in the absence of cytoplasmic enzyme secretion, after phagocytosis of chrysotile.

It has been demonstrated that macrophages respond differently to different fibers and doses and that the response is also dependent on the animal species used (6-8). However, it is known that the macrophage is an immunocompetent cell which performs a number of functions connected with the defense mechanisms of the lung. Although these im-

munological functions have only been poorly examined after fiber exposures (9), the production of lysozyme and a phagocytic ability are known to be among its capabilities.

The geometrical dimensions of fibers seem to be an important factor in the pathogenesis of both fibrosis and cancer.

Materials and Methods

Cell Cultures of Alveolar Macrophages

Unstimulated guinea pig and rat lung macrophages were obtained by pulmonary lavage using heparinized (8 U/mL) calcium- and magnesium-free phosphate-buffered saline (PBS) (10, 11). The resulting cell suspension was washed three times in heparin-free PBS, centrifuged at 350g, resuspended in serum-free Minimal Essential Medium (MEM) and adjusted to 1×10^6 cells/mL. Rat lung macrophages, 0.1×10^6 , were then transferred to the cavities of microtiter plates (Falcon Plastics, No. 3040F) and 0.75×10^6 guinea pig cells were transferred to the cavities of multiwell plates (Falcon Plastics, No. 3008).

After 2 hr incubation at 37°C in a 5% CO₂/95%

*Hygiene-Institute of Justus Liebig University, Giessen, Federal Republic of Germany.

air mixture to allow cell attachment, the MEM was removed and different doses of the dusts were added in McCoy's 5A medium in the presence or absence of heat inactivated fetal calf serum.

For phagocytic determination, the cells were cultured in special glass vessels at a concentration of 0.2×10^6 cells/tube, but the incubation conditions were the same as those given above.

Fibers and Control Dusts

Fiber fractions from different sources were used. Some of them were prepared by using the technique of Spurny et al. (12) as previously described (13). The other fibers were a gift of The Johns Manville Corp (U.S.A.), and their physical characteristics are shown in Table 1.

Table 1. Length and diameter distributions of fibers.

No.	Fiber type	Length distribution		Diameter distribution	
		Length, μm	% of fibers	Diameter, μm	% of fibers
1	Short, very thin glass fibers (4106-23-7)	<1	83.8	<0.1	52.0
		1.0-1.9	11.5	0.1-0.14	35.4
		>1.9	4.8	0.15-0.19	5.7
				0.2-0.29	6.9
2	Long, very thin glass fibers (4106-24-2)	1.0-2.9	9.2	<0.1	63.7
		3.0-4.9	15.8	0.1-0.14	25.6
		5.0-9.9	25.0	0.15-0.19	8.8
		10-19	30.0	0.2-0.29	1.6
		20-29	9.2	0.3-0.59	0.3
		30-59	10.8	(0.41)	
3	Short, thin glass fibers (4106-4-3)	1.0-1.9	14.1	<0.15	5.9
		2.0-2.9	35.2	0.15-0.29	16.4
		3.0-4.9	35.2	0.3-0.59	35.4
		5.0-9.9	8.5	0.6-0.99	26.2
		10-22	7.0	1.0-1.99	16.1
4	Long, thin glass fibers (4106-19-2)	<5	0.7	<0.15	51.4
		5.9-9	7.0	0.15-0.19	19.6
		10-19	37.5	0.20-0.29	11.9
		20-39	36.0	0.30-0.39	6.1
		40-59	14.8	0.40-0.59	6.2
		60-99	4.0	0.60-1.49	4.8
5	Short, thick glass fibers (4106-6-1)	1.0-2.9	22.4	0.4-0.59	3.4
		3.0-4.9	33.5	0.6-0.99	5.8
		5.0-9.9	31.5	1.00-1.49	15.2
		10-19	10.5	1.50-1.99	28.6
		20-29	2.1	3.00-3.95	17.6
6	Long, thick glass fibers (4106-19-1a)	3.0-4.9	9.9	0.2-0.59	4.1
		5.0-9.9	14.6	0.6-0.99	18.0
		10-19	28.9	1.0-1.49	35.2
		20-29	20.1	1.5-1.99	22.7
		30-59	22.1	2.0-2.95	14.8
		60-99	4.4	3.0-5.95	5.2
7	Jeffrey fibrils (chrysotile) (4173-46-1)	<10	29.5	0.03-0.05	99.9
		1.0-1.9	34.7		
		2.0-4.9	27.6		
		5.0-9.9	6.6		
		10-19	1.6		
8	Jeffrey fibrils (chrysotile) (4173-46-2)	0.10-0.99	11.8	0.10	99.9
		1.0-1.9	20.0		
		2.0-4.9	24.1		
		5.0-9.9	20.5		
		10-19	17.7		
9	Short Munroe fibrils (chrysotile) (4173-60-1)	0.40-9.99	22.5	0.03-0.05	99.9
		1.0-1.49	29.7		
		1.5-1.99	21.0		
		2.0-2.99	15.3		
		3.0-4.99	10.0		
	5.0-9.99	1.7			

Table 1. (Continued).

No.	Fiber type	Length distribution		Diameter distribution	
		Length, μm	% of fibers	Diameter, μm	% of fibers
10	Long Munroe chrysotile (4173-60-2)	<3.0	0.0	<0.1	40.2
		3.0-4.9	0.7	0.1-0.29	20.2
		5.0-9.9	4.3	0.3-0.59	8.8
		10-19	26.4	0.6-0.99	13.6
		20-60	65.7	1.0-1.99	4.2
11	Short crocidolite (S. Africa)	60-80	2.9	2.0-2.5	13.3
		0.40-0.99	17.6	0.03-0.04	1.2
		1.0-1.9	44.4	0.05-0.09	7.5
		2.0-4.9	19.8	0.10-0.19	43.3
		5.0-9.9	6.2	0.20-0.39	32.4
12	Short crocidolite (4106-23-9)	10-19	1.6	0.40-0.59	4.4
		20-29	0.4	0.60-0.99	1.2
		0.1-0.69	47.8	0.05-0.09	25.3
		0.7-0.99	22.3	0.10-0.14	40.0
		1.0-2.9	24.6	0.15-0.19	19.4
13	Long crocidolite (4106-22-4)	3.0-4.9	3.5	0.20-0.29	13.3
		5.0-9.9	1.5	0.30-0.39	2.0
		10-19	0.3		
		3.0-4.9	4.2	0.10-0.14	27.8
		5.0-9.9	13.6	0.15-0.19	15.0
14	Short fluoroamphibole (4106-31-3)	10-19	28.3	0.20-0.29	30.9
		20-29	23.0	0.30-0.39	13.1
		30-59	24.1	0.4-1.49	7.9
		60-129	6.8		
		1	89.9	0.03-0.09	46.0
15	Respirable fluoroamphibole (4106-30-1)	1.0-1.9	5.8	0.10-0.19	40.9
		2.0-4.9	3.6	0.20-0.39	9.6
		5.0-19.0	0.75	0.40-0.99	2.9
		1.0-2.95		1.0-2.95	0.6
		0.5-1.49	8.2	0.03-0.04	44.7
		1.5-2.9	21.5	0.05-0.14	25.3
		3.0-4.9	27.0	0.15-0.39	20.9
		5.0-9.9	27.5	0.40-0.99	5.6
		10-19	12.5	1.0-2.0	3.5
		20-39	3.5		

The samples prepared by using the technique of Spurny et al. (12) have been defined by length, but those from The Johns Manville Corp. have also been defined by diameter.

Dörentruper quartz (DQ₁₂) and electrocorundum with a similar size distribution served as control dusts.

All of the dust samples were weighed and sterilized by dry heat before being suspended in medium by the use of ultrasonics.

Assay Systems

Lactate dehydrogenase (LDH) and lactic acid were estimated by using biochemical test kits (Boehringer GmbH, Mannheim, FRG). β -Glucuronidase was also measured by using a test kit (Sigma Chemie GmbH, München, FRG) and the lysozyme concentration was measured by using reagents supplied by Behringwerke AG (Marburg, FRG).

Phagocytosis was assayed quantitatively by de-

termining the amount of luminescence produced after the addition of serum-opsonized zymosan A particles (Sigma Chemie GmbH, München, FRG). The chemiluminescence was measured after luminol (Lumac, Düsseldorf, FRG) enhancement in a six-channel apparatus (Biolumat 9505; Berthold, Wildbad, FRG). The results were printed out in counts/minute (cpm) using a teletype.

Results and Discussion

Modified UICC samples, well-defined fiber fractions and also glass fibers (12) cause release of the cytoplasmic enzyme LDH and the lysosomal enzyme β -glucuronidase from guinea pig alveolar macrophages after incubation with the relatively low concentration of 100 fibers/cell (Table 2). At this concentration, we do not find a significant difference between the two parameters. Table 2 also shows that the cytotoxicity is dependent on fiber length for all four fiber types. It is interesting to note that

Table 2. Enzyme release of guinea pig lung macrophages 24 hr after incubation with asbestos and glass fiber fractions of three well-defined length spectra.^a

Expt. no.		Enzyme release, mU/mL	
		LDH	β -Glucuronidase
1	K _o	31	58
	K _{rd}	35	53
	K _F	112	98
	K _{FF}	68	101
	K _{fF}	47	45
2	K _o	30	52
	K _{rd}	31	49
	A _F	92	90
	A _{FF}	60	75
	A _{fF}	61	74
3	K _o	23	33
	K _{rd}	26	35
	C _f	51	83
	C _{FF}	44	57
	C _{fF}	41	48
4	K _o	28	37
	K _{rd}	31	34
	G _{F100}	88	74
	G _{FF}	70	61
	G _{fF}	42	40

^a Dust concentrations = 100 fibers/cell.

chrysotile showed a relatively low toxicity in comparison with crocidolite, amosite and glass fiber. This may be due to the substantial gravimetric differences that occur when the same numerical dose is used. For example, there would be a 2-fold weight increase for crocidolite and a 10-fold increase for amosite and glass fiber when compared to the weight of chrysotile.

These findings are in agreement with the results of Davies (8) and demonstrate that, at low fiber concentrations, any differential release of β -glucuronidase relative to LDH is low in contrast to that seen at higher concentrations (6, 7).

Figures 1-3 show the results obtained with 15 new fiber fractions of glass fiber, chrysotile, crocidolite and synthetic fluoroamphiboles, all of which had interesting geometric dimensions. Figure 1 shows the amount of LDH released into the medium in the presence of serum after an incubation time of 20 hr. The results are expressed as a percentage with the cell control equaling 0% and quartz DQ₁₂ equaling 100%. Figure 2 gives the equivalent results obtained when serum was absent and Figure 3 shows the lactic acid production, after a similar incubation period in the absence of serum, and the results are expressed as a percentage of the lactic acid produced by the control cells. With the exception of the synthetic amphibole, LDH release was dependent on fiber length in the presence or absence of serum in the medium.

A comparison of the three pairs of glass fiber




FIGURE 1. LDH release of guinea pig lung macrophages 20 hr after incubation with 100 μ g dust/ 10^6 cells. Control = 0%; DQ₁₂ = 100%.




FIGURE 2. LDH release of guinea pig lung macrophages 20 hr after incubation with 100 μ g dust/ 10^6 cells. Control = 0%; DQ₁₂ = 100%.

samples shows that fiber toxicity is clearly dependent on both diameter and length. As no data were available on the number of fibers per unit weight, it was not possible to do further tests with different numerical concentrations. In contrast to our earlier investigations (13), it can be seen that even every short fiber can exhibit a high toxicity. The important difference between this study and the earlier tests lies in the very thin fiber diameter (thinner than 0.03 μ m) in this study, in comparison to a mean diameter of 0.19 μ m in the preparation made by Spurny et al. (12). In fiber fractions with a diameter of 1.0 μ m, only the longer fibers cause a high release of LDH. Short fibers of this diameter and both fiber




FIGURE 3. Lactic acid production of guinea pig lung macrophages 20 hr after incubation with $100 \mu\text{g}$ dust/ 10^6 cells. Control = 100%; DQ_{12} = 0%.




FIGURE 4. Lysozyme production of guinea pig lung macrophages 72 hr after incubation with electrocorundum and UICC asbestos at a dust concentration of $100 \mu\text{g}/10^6$ cells: (A) control; (B) electrocorundum; (C) UICC chrysotile; (D) UICC crocidolite; (E) UICC amosite.




FIGURE 5. Phagocytosis determined by bioluminescence with opsonized zymosan following luminol enhancement 18 hr after incubation of guinea pig lung macrophages with asbestos and glass fibers. Graph shows one determination in counts per minute (cpm).

samples with a diameter of $3 \mu\text{m}$ are not toxic in our system.

A comparison of the results obtained when the cells were incubated in the presence or absence of serum gave no additional information on fiber toxicity. Glass fiber caused a considerably lower reduction in lactic acid production in comparison to the chrysotile and crocidolite fiber fractions with similar dimensions. Possibly, the surface properties of the different fibers play a greater role here than in causing the release of cytoplasmic enzymes. The influence of the UICC asbestos samples on lysozyme production was small and was only detected at high concentrations (Fig. 4). At very low concentrations we found an enhancement of cellular activity.

Figure 5 demonstrates the phagocytic activity of alveolar macrophages after 20 hr exposure to UICC asbestos samples and JM 100 glass fiber. Chemiluminescence generated by the phagocytic process was measured after Luminol enhancement. Using this assay it was shown that concentrations of $100 \mu\text{g}$ dust/mL caused a significant depression of phagocytosis for all four fiber types. Due to technical problems, only some of the results from this assay could be evaluated. In the future, we plan to investigate the effect of lower dust concentrations and longer exposure times on this test system in order to study the direct influence of dust loading on the phagocytic ability of the cells.

In conclusion, we wish to stress that more fiber fractions with narrowly defined length and diameter spectra are needed in order to make better and more meaningful comparisons between fibers with different chemical compositions.

REFERENCES

- Davies, P., and Allison, A. C. Secretion of macrophage enzymes in relation to the pathogenesis of chronic inflammation. In: *Immunobiology of the Macrophage* (D. S. Nelson, Ed.), Academic Press, New York, 1976, pp. 427-461.
- Loose, L. D., Silkworth, J. B., Charbonneau, T., and Blumenstock, F. Environmental chemical-induced macrophage dysfunction. *Environ. Health Perspect.* 39: 79-91 (1981).
- Allison, A. C., Harington, J. S., and Bierbeck, M. An examination of the cytotoxic effects of silica on macrophages. *J. Exptl. Med.* 124: 141-154 (1966).
- Beck, E. G., Holt, P. F. and Manojlovic, N. Comparison of effects on macrophage cultures of glass fibre, glass powder and chrysotile asbestos. *Brit. J. Ind. Med.* 29: 280-286 (1972).
- Davies P., Allison, A. C., Ackermann, J. Butterfield, A., and Williams S. Asbestos induces selective release of lysosomal enzymes from mononuclear phagocytes. *Nature* 251: 423-425 (1974).
- Morgan, D. M. L. and Allison, A. C. Effects of silica and asbestos on alveolar and peritoneal macrophages: a com-

- parative study. In: *The In Vitro Effects of Mineral Dusts* (R. C. Brown, I. P. Gormley, M. Chamberlain and R. Davies, Eds.), Academic Press, London, 1980, pp. 75-81.
- Jaurand, M. C., Magne, L., and Bignon, J. Effects well-defined fibres on red blood cells and alveolar macrophages. In: *Biological Effects of Mineral Fibres* IARC Sci. Publ. 1(30) (J.C. Wagner, Ed.), International Agency for Research on Cancer, Lyon, 1980, pp. 441-450.
 - Davies, R. The effect of dusts on enzyme release from macrophages. In: *The In Vitro Effects of Mineral Dusts* (R. C. Brown, I. P. Gormley, M. Chamberlain, and R. Davies, Eds.), Academic Press, London, 1980. pp. 67-74.
 - Miller, K. Alteration in the surface-related phenomena of alveolar macrophage following inhalation of crocidolite asbestos and quartz dusts: and overview. *Environ. Res.* 20: 162-182 (1979).
 - Myrik, Q. N., Leake, E. S., and Oshima, S. A study of macrophages and epithelioid-like cells from granulomatous (BCG-induced) lungs of rabbit. *J. Immunology* 85: 745-752 (1962).
 - Beck E. G. Die Reaktion in vitro gezüchteter Zellen auf partikelförmige Luftverunreinigungen und hochpolymere Stoffe. *Forschungsber. Landes Nordrhein-Westfalen* 2083: 1-125 (1970).
 - Spurny, K., Stober, V., Opiela, H., and Weiss, G. Size-selective preparation of inorganic fibres for biological experiments. *Am. Ind. Hyg. Assoc. J.* 40: 20-38 (1979).
 - Tilkes, F., and Beck, E. G. Comparison of length-dependent cytotoxicity of inhalable asbestos and man-made mineral fibres. In: *Biological Effects of Mineral Fibres*, IARC Sci. Publ. 1(30) (J. C. Wagner, Ed.), International Agency for Research on Cancer, Lyon 1980, pp. 475-483.